

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **CD20 / MS4A1 (B-Cell Marker) (IGEL/1497R), 0.2mg/mL, Clone: [IGEL/1497R], Rabbit, Monoclonal BOT-BNUB1497-500**

|                            |   |
|----------------------------|---|
| Article Name               | CD20 / MS4A1 (B-Cell Marker) (IGEL/1497R), 0.2mg/mL, Clone: [IGEL/1497R], Rabbit, Monoclonal  |
| Biozol Catalog Number      | BOT-BNUB1497-500  |
| Supplier Catalog Number    | BNUB1497-500  |
| Alternative Catalog Number | BOT-BNUB1497-500-500UL  |
| Manufacturer               | Biotium   |
| Host                       | Rabbit  |
| Category                   | Antikörper  |
| Application                | FC, IHC   |
| Species Reactivity         | Human   |
| Immunogen                  | Recombinant human full-length MS4A1 protein   |
| Product Description        | This antibody recognizes a protein of 30-33 kDa, which is identified as CD20. It is a non-Ig differentiation antigen of B-cells and its expression is restricted to normal and neoplastic B-cells, being absent from all other leukocytes and tissues. CD2... |
| Clonality                  | Monoclonal  |
| Concentration              | 0.2 mg/mL   |
| Clone Designation          | [IGEL/1497R]  |
| Molecular Weight           | 33-37 kDa   |

|                   |  |
|-------------------|--|
| UniProt           | <a href="#">P11836</a>   |
| Buffer            | PBS, 0.05% BSA, 0.05% azide  |
| Source            | Animal   |
| Application Notes | Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Immunofluorescence 0.5-1 ug/mL Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |