

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### CD162 (Selectin P Ligand) (PSGL1/1601), 0.2mg/mL, IgG1, Clone: [PSGL1/1601], Mouse, Monoclonal BOT-BNUB1601-100

Article Name	CD162 (Selectin P Ligand) (PSGL1/1601), 0.2mg/mL, IgG1, Clone: [PSGL1/1601], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUB1601-100
Supplier Catalog Number	BNUB1601-100
Alternative Catalog Number	BOT-BNUB1601-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant human full-length SELPLG (CD162) protein
Product Description	CD162 glycoprotein functions as a high affinity counter-receptor for the cell adhesion molecules P-, E- and L- selectin expressed on myeloid cells and stimulated T lymphocytes. As such, this protein plays a critical role in leukocyte trafficking duri...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[PSGL1/1601]
Molecular Weight	43-120 kDa

Isotype	IgG1
UniProt	<a href="#">Q14242</a>
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	<p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p>