

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **CD56 / NCAM1 / NKH1 (Neuronal Cell Marker)(ERIC-1), 0.2mg/mL, Clone: [ERIC-1], Mouse, Monoclonal BOT-BNUB1602-500**

Article Name	CD56 / NCAM1 / NKH1 (Neuronal Cell Marker)(ERIC-1), 0.2mg/mL, Clone: [ERIC-1], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUB1602-500
Supplier Catalog Number	BNUB1602-500
Alternative Catalog Number	BOT-BNUB1602-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human, Rat
Immunogen	Recombinant human NCAM1 protein
Product Description	This MAbs reacts with an extracellular domain (close to transmembrane) of CD56/NCAM. Three isoforms of neural cell adhesion molecule (NCAM) are produced by differential splicing of the RNA transcript from a single gene. The 135 kDa isoform is the basi...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[ERIC-1]
Molecular Weight	180, 145 and 125 kDa

UniProt	P13591
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	<p>For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry</p>