

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Chromogranin A / CHGA (Neuroendocrine Marker) (CHGA/1731R), 0.2mg/mL, Clone: [CHGA/1731R], Rabbit, Monoclonal BOT-BNUB1731-500

| | |
|----------------------------|---|
| Article Name | Chromogranin A / CHGA (Neuroendocrine Marker) (CHGA/1731R), 0.2mg/mL, Clone: [CHGA/1731R], Rabbit, Monoclonal |
| Biozol Catalog Number | BOT-BNUB1731-500 |
| Supplier Catalog Number | BNUB1731-500 |
| Alternative Catalog Number | BOT-BNUB1731-500-500UL |
| Manufacturer | Biotium |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant human full-length Chromogranin A protein |
| Product Description | Chromogranin A is present in neuroendocrine cells throughout the body, including the neuroendocrine cells of the large and small intestine, adrenal medulla and pancreatic islets. It is an excellent marker for carcinoid tumors, pheochromocytomas, para... |
| Clonality | Monoclonal |
| Concentration | 0.2 mg/mL |
| Clone Designation | [CHGA/1731R] |
| Molecular Weight | 68-75 kDa |

| | |
|-------------------|--|
| UniProt | P10645 |
| Buffer | PBS, 0.05% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |