

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker) (VWF/1767), 0.2mg/mL, Clone: [VWF/1767], Mouse, Monoclonal BOT-BNUB1767-500

Article Name	von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker) (VWF/1767), 0.2mg/mL, Clone: [VWF/1767], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUB1767-500
Supplier Catalog Number	BNUB1767-500
Alternative Catalog Number	BOT-BNUB1767-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant fragment of human vWF protein (aa1815-1939) (exact sequence is proprietary)
Product Description	von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[VWF/1767]
Molecular Weight	250 kDa

UniProt	P04275
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody ELISA: 2-4 ug/mL for coating order Ab without BSA Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user