

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD45RB (B-Cell Marker) (PTPRC/1783R), 0.2mg/mL, Clone: [PTPRC/1783R], Rabbit, Monoclonal BOT-BNUB1783-100

| | |
|----------------------------|---|
| Article Name | CD45RB (B-Cell Marker) (PTPRC/1783R), 0.2mg/mL, Clone: [PTPRC/1783R], Rabbit, Monoclonal |
| Biozol Catalog Number | BOT-BNUB1783-100 |
| Supplier Catalog Number | BNUB1783-100 |
| Alternative Catalog Number | BOT-BNUB1783-100-100UL |
| Manufacturer | Biotium |
| Host | Rabbit |
| Category | Antikörper |
| Application | FC, IF, IHC, WB |
| Species Reactivity | Human |
| Immunogen | Human B-lymphocytes |
| Product Description | CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms ar... |
| Clonality | Monoclonal |
| Concentration | 0.2 mg/mL |
| Clone Designation | [PTPRC/1783R] |
| Molecular Weight | 180-220 kDa |

| | |
|-------------------|---|
| UniProt | P08575 |
| Buffer | PBS, 0.05% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user</p> |