

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Histone H1 (Nuclear Marker) (HH1/1784R), 0.2mg/mL, Clone: [HH1/1784R], Rabbit, Monoclonal BOT-BNUB1784-100

Article Name	Histone H1 (Nuclear Marker) (HH1/1784R), 0.2mg/mL, Clone: [HH1/1784R], Rabbit, Monoclonal
Biozol Catalog Number	BOT-BNUB1784-100
Supplier Catalog Number	BNUB1784-100
Alternative Catalog Number	BOT-BNUB1784-100-100UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Application	FC, IF, IHC, WB
Species Reactivity	Human, Mouse, Rat
Immunogen	Recombinant full-length human Histone H1 protein
Product Description	Eukaryotic histones are basic and water-soluble nuclear proteins that form hetero-octameric nucleosome particles by wrapping 146 base pairs of DNA in a left-handed superhelical turn sequentially to form chromosomal fiber. Two molecules of each of...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[HH1/1784R]
Molecular Weight	~30 kDa

UniProt	Multiple
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user