

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Ep-CAM / CD326 (Epithelial Marker) (rVU-1D9), 0.2mg/mL, Clone: [rVU-1D9], Mouse, Monoclonal BOT-BNUB1820-100**

Article Name	Ep-CAM / CD326 (Epithelial Marker) (rVU-1D9), 0.2mg/mL, Clone: [rVU-1D9], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUB1820-100
Supplier Catalog Number	BNUB1820-100
Alternative Catalog Number	BOT-BNUB1820-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	Exosome Staining, FC, IF, IHC, WB
Species Reactivity	Human
Immunogen	Small cell lung carcinoma cells
Product Description	This antibody reacts with the first EGF repeat in the extracellular domain of Ep-CAM. It is a 40-43 kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[rVU-1D9]
Molecular Weight	40-43 kDa

UniProt	<a href="#">P16422</a>
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user