

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD30 / TNFRSF8 (Hodgkin & Reed-Sternberg Cell Marker) (rKi-1/779), 0.2mg/mL, Clone: [rKi-1/779], Mouse, Monoclonal BOT-BNUB1828-100

| | |
|----------------------------|---|
| Article Name | CD30 / TNFRSF8 (Hodgkin & Reed-Sternberg Cell Marker) (rKi-1/779), 0.2mg/mL, Clone: [rKi-1/779], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNUB1828-100 |
| Supplier Catalog Number | BNUB1828-100 |
| Alternative Catalog Number | BOT-BNUB1828-100-100UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant full-length human TNFRSF8 protein |
| Product Description | This antibody recognizes a single chain glycoprotein of 105/120 kDa, identified as CD30/Ki-1. CD30 is synthesized as a 90 kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120 kDa glycoprotein. In ... |
| Clonality | Monoclonal |
| Concentration | 0.2 mg/mL |
| Clone Designation | [rKi-1/779] |
| Molecular Weight | 105-120 kDa |

| | |
|-------------------|---|
| UniProt | P28908 |
| Buffer | PBS, 0.05% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 1 mM EDTA pH 7.5-8.5 Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |