

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Wilms Tumor 1 (WT1) (Wilms Tumor & Mesothelial Marker) (rWT1/857), 0.2mg/mL, Clone: [rWT1/857], Mouse, Monoclonal BOT-BNUB1856-500

Article Name	Wilms Tumor 1 (WT1) (Wilms Tumor & Mesothelial Marker) (rWT1/857), 0.2mg/mL, Clone: [rWT1/857], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUB1856-500
Supplier Catalog Number	BNUB1856-500
Alternative Catalog Number	BOT-BNUB1856-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant full-length human WT1 protein
Product Description	This antibody recognizes a 47-55 kDa-tumor suppressor protein, identified as Wilms Tumor (WT1) protein. The antibody reacts with all isoforms of the full-length WT1 and also identifies WT1 lacking exon 2-encoded amino acids, frequently found in subse...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[rWT1/857]
Molecular Weight	47-55 kDa

UniProt	P19544
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user