

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Beta-2 Microglobulin (Renal Failure & Tumor Marker) (B2M/1857R), 0.2mg/mL, Clone: [B2M/1857R], Rabbit, Monoclonal BOT-BNUB1857-100**

Article Name	Beta-2 Microglobulin (Renal Failure & Tumor Marker) (B2M/1857R), 0.2mg/mL, Clone: [B2M/1857R], Rabbit, Monoclonal
Biozol Catalog Number	BOT-BNUB1857-100
Supplier Catalog Number	BNUB1857-100
Alternative Catalog Number	BOT-BNUB1857-100-100UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Application	FC, IF, IHC, WB
Species Reactivity	Human, Primate
Immunogen	Recombinant full-length human B2M protein
Product Description	Beta-2 microglobulin is a component of the class I major histocompatibility complex (MHC I) where it functions in antigen processing and presentation. Involved in the presentation of peptide antigens to the immune system. Human Beta-2 microglobulin i...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[B2M/1857R]
Molecular Weight	12 kDa

UniProt	<a href="#">P61769</a>
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user