

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### CD27 (Tumor Necrosis Factor Receptor Superfamily 7) (LPFS2/2034R), 0.2mg/mL, Clone: [LPFS2/2034R], Rabbit, Monoclonal BOT-BNUB2034-100

Article Name	CD27 (Tumor Necrosis Factor Receptor Superfamily 7) (LPFS2/2034R), 0.2mg/mL, Clone: [LPFS2/2034R], Rabbit, Monoclonal
Biozol Catalog Number	BOT-BNUB2034-100
Supplier Catalog Number	BNUB2034-100
Alternative Catalog Number	BOT-BNUB2034-100-100UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant full-length human CD22 protein (exact sequence is proprietary)
Product Description	This antibody recognizes a protein of a disulfide-linked 120 kDa dimer, identified as CD27. It is expressed on the majority of peripheral T cells, medullary thymocytes, memory-type B cells, and natural killer cells. It is a transmembrane phosphoglyco...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[LPFS2/2034R]
Molecular Weight	120 kDa

UniProt	P26842
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris buffer with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user