

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Cytokeratin 16 (KRT16) (Suprabasal Keratinocyte Marker) (KRT16/2043R), 0.2mg/mL, Clone: [KRT16/2043R], Rabbit, Monoclonal BOT-BNUB2043-100

Article Name	Cytokeratin 16 (KRT16) (Suprabasal Keratinocyte Marker) (KRT16/2043R), 0.2mg/mL, Clone: [KRT16/2043R], Rabbit, Monoclonal
Biozol Catalog Number	BOT-BNUB2043-100
Supplier Catalog Number	BNUB2043-100
Alternative Catalog Number	BOT-BNUB2043-100-100UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant fragment from the C-terminal of human Cytokeratin 16
Product Description	Cytokeratins are a family of intermediate filament proteins that assemble into filaments through forming heterodimers of one type I Cytokeratins (Cytokeratins 9 to 23) and one type II Cytokeratins (keratins 1 to 8). The cytokeratin proteins play a cr...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[KRT16/2043R]
Molecular Weight	48 kDa

UniProt	P08779
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user