

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

TNFS15 / VEGI (Vascular Endothelial Growth Inhibitor) (VEGI/2052R), 0.2mg/mL, Clone: [VEGI/2052R], Rabbit, Monoclonal BOT-BNUB2052-500

| | |
|----------------------------|---|
| Article Name | TNFS15 / VEGI (Vascular Endothelial Growth Inhibitor) (VEGI/2052R), 0.2mg/mL, Clone: [VEGI/2052R], Rabbit, Monoclonal |
| Biozol Catalog Number | BOT-BNUB2052-500 |
| Supplier Catalog Number | BNUB2052-500 |
| Alternative Catalog Number | BOT-BNUB2052-500-500UL |
| Manufacturer | Biotium |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant full-length human VEGI protein |
| Product Description | VEGI is an anti-angiogenic cytokine that belongs to tumor necrosis factor superfamily, member 15 (TNFSF15). This protein is abundantly expressed in endothelial cells, but is not expressed in either B or T cells. The expression of this protein is indu... |
| Clonality | Monoclonal |
| Concentration | 0.2 mg/mL |
| Clone Designation | [VEGI/2052R] |
| Molecular Weight | 22 kDa |

| | |
|-------------------|--|
| UniProt | O95150 |
| Buffer | PBS, 0.05% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |