

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Bcl-6 (Follicular Lymphoma Marker) (rBCL6/1718), 0.2mg/mL, Clone: [rBCL6/1718], Mouse, Monoclonal BOT-BNUB2498-100

Article Name	Bcl-6 (Follicular Lymphoma Marker) (rBCL6/1718), 0.2mg/mL, Clone: [rBCL6/1718], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUB2498-100
Supplier Catalog Number	BNUB2498-100
Alternative Catalog Number	BOT-BNUB2498-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant human bcl-6 protein fragment (aa256-389) (exact sequence is proprietary)
Product Description	Bcl-6 is a transcriptional repressor mainly required for germinal center (GC) formation and antibody affinity maturation which has different mechanisms of action specific to the lineage and biological functions. It forms complexes with different core...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[rBCL6/1718]
Molecular Weight	95 kDa

UniProt	P41182
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user