

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

TCL1 (T-Cell Marker) (TCL1/2747R), 0.2mg/mL, Clone: [TCL1/2747R], Rabbit, Monoclonal BOT-BNUB2747-100

Article Name	TCL1 (T-Cell Marker) (TCL1/2747R), 0.2mg/mL, Clone: [TCL1/2747R], Rabbit, Monoclonal
Biozol Catalog Number	BOT-BNUB2747-100
Supplier Catalog Number	BNUB2747-100
Alternative Catalog Number	BOT-BNUB2747-100-100UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant human TCL1 protein fragment (around aa 2-109) (exact sequence is proprietary)
Product Description	T-cell leukemia/Lymphoma Protein 1A (TCL-1A), also known as p14TCL1 is a product of the TCL1 gene that is involved in T-cell prolymphocytic leukemia (TPLL). T-PLL is a rare form of mature T-cell leukemia, which is consistently associated with chromos...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[TCL1/2747R]
Molecular Weight	14 kDa

UniProt	P56279
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user