

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

AMACR / p504S (Prostate Cancer Marker) (AMACR/2748R), 0.2mg/mL, Clone: [AMACR/2748R], Rabbit, Monoclonal BOT-BNUB2748-500

Article Name	AMACR / p504S (Prostate Cancer Marker) (AMACR/2748R), 0.2mg/mL, Clone: [AMACR/2748R], Rabbit, Monoclonal
Biozol Catalog Number	BOT-BNUB2748-500
Supplier Catalog Number	BNUB2748-500
Alternative Catalog Number	BOT-BNUB2748-500-500UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Application	IHC, WB
Species Reactivity	Human
Immunogen	Recombinant human AMACR protein fragment (around aa 297-394) (exact sequence is proprietary)
Product Description	This antibody recognizes a protein of 42 kDa, which is identified as Alpha-methylacyl-CoA Racemase (AMACR), also known as p504S. It is an enzyme that is involved in bile acid biosynthesis and betbeta-oxidation of branched-chain fatty acids. AMACR is ...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[AMACR/2748R]
Molecular Weight	42 kDa

UniProt	Q9UHK6
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user