

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Golgi Complex (Marker for Human Cells) (GLG1/2829R), 0.2mg/mL, Clone: [GLG1/2829R], Rabbit, Monoclonal BOT-BNUB2829-100

Article Name	Golgi Complex (Marker for Human Cells) (GLG1/2829R), 0.2mg/mL, Clone: [GLG1/2829R], Rabbit, Monoclonal
Biozol Catalog Number	BOT-BNUB2829-100
Supplier Catalog Number	BNUB2829-100
Alternative Catalog Number	BOT-BNUB2829-100-100UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	SU-DHL-1 large cell lymphoma cells.
Product Description	This MAb recognizes an antigen associated with the Golgi complex in human cells only. It can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining ...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[GLG1/2829R]
Molecular Weight	Not Known

UniProt	Not Known
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Does not react with mouse or rat Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user