

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD79b (B-Cell Marker)(IGB/2940R), 0.2mg/mL, Clone: [IGB/2940R], Rabbit, Monoclonal BOT-BNUB2940-100

Article Name	CD79b (B-Cell Marker)(IGB/2940R), 0.2mg/mL, Clone: [IGB/2940R], Rabbit, Monoclonal
Biozol Catalog Number	BOT-BNUB2940-100
Supplier Catalog Number	BNUB2940-100
Alternative Catalog Number	BOT-BNUB2940-100-100UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Species Reactivity	Human
Immunogen	Recombinant human CD79b protein fragment (around aa29-159) (exact sequence is proprietary)
Product Description	CD79 (also designated Ig chains, designated CD79B or B29. The B cell antigen receptor complex (BCR) is formed by the association of CD79 with a membrane immunoglobulin, such as IgM or IgD. The membrane immunoglobulins IgM and IgD achieve surface expr...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[IGB/2940R]
Molecular Weight	39 kDa
UniProt	P40259

Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry