

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

HLA-Pan (MHC II) (HLA-Pan/2967R), 0.2mg/mL, Clone: [HLA-Pan/2967R], Rabbit, Monoclonal BOT-BNUB2967-100

| | |
|----------------------------|---|
| Article Name | HLA-Pan (MHC II) (HLA-Pan/2967R), 0.2mg/mL, Clone: [HLA-Pan/2967R], Rabbit, Monoclonal |
| Biozol Catalog Number | BOT-BNUB2967-100 |
| Supplier Catalog Number | BNUB2967-100 |
| Alternative Catalog Number | BOT-BNUB2967-100-100UL |
| Manufacturer | Biotium |
| Host | Rabbit |
| Category | Antikörper |
| Application | FC, IF, IHC, WB |
| Species Reactivity | Human |
| Immunogen | Non-T, non-B human acute lymphoblastic leukemia REH6 cell line |
| Product Description | This antibody reacts with a common epitope of human major histocompatibility (MHC) class II antigens, HLA-DP, -DQ and -DR. Human MHC class II antigens are transmembrane glycoproteins composed of an alpha chain (36 kDa) and a beta chain (27 kDa). They... |
| Clonality | Monoclonal |
| Concentration | 0.2 mg/mL |
| Clone Designation | [HLA-Pan/2967R] |
| Molecular Weight | 33-35 kDa |

| | |
|-------------------|--|
| UniProt | P04440 |
| Buffer | PBS, 0.05% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody ELISA: For coating use Ab at 1-5 ug/mL order Ab without BSA Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes |