

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

BCL10 (MALT-Lymphoma Marker) (BL10/2988R), 0.2mg/mL, Clone: [BL10/2988R], Rabbit, Monoclonal BOT-BNUB2988-500

Article Name	BCL10 (MALT-Lymphoma Marker) (BL10/2988R), 0.2mg/mL, Clone: [BL10/2988R], Rabbit, Monoclonal
Biozol Catalog Number	BOT-BNUB2988-500
Supplier Catalog Number	BNUB2988-500
Alternative Catalog Number	BOT-BNUB2988-500-500UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Application	FC, IF, IHC
Species Reactivity	Human
Immunogen	Recombinant human BCL10 protein fragment (around aa122-168) (exact sequence is proprietary)
Product Description	B-cell lymphoma/leukemia 10 (BCL10) is a signaling protein that promotes apoptosis, pro-caspase-9 maturation, and activation of NF-kappa-B via NIK and IKK. BCL10 contains an N-terminal caspase recruitment domain (CARD), which is found in a number of ...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[BL10/2988R]
Molecular Weight	33 kDa

UniProt	O95999
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user