

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Nucleolar / Nucleoli(NM95), 1mg/mL, Clone: [NM95], Mouse, Monoclonal BOT-BNUM0095-50

| | |
|----------------------------|---|
| Article Name | Nucleolar / Nucleoli(NM95), 1mg/mL, Clone: [NM95], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNUM0095-50 |
| Supplier Catalog Number | BNUM0095-50 |
| Alternative Catalog Number | BOT-BNUM0095-50-50UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | FC, IF, IHC |
| Species Reactivity | Human |
| Immunogen | Nuclei of myeloid leukemia biopsy cells |
| Product Description | This MAb is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells. This monoclonal antibody is part of a new panel of reagents, which recognizes subcellular organelles or compartments of human cell... |
| Clonality | Monoclonal |
| Concentration | 1 mg/mL |
| Clone Designation | [NM95] |
| Molecular Weight | Not Known |
| UniProt | Not Known |

| | |
|-------------------|--|
| Buffer | PBS, no BSA, no azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Does not react with mouse, rat or cow, others not known Immunohistology formalin-paraffin 0.25-0.5 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Immunocytochemistry acetone-fixed cells 0.5-1.0 ug/mL Immunofluorescence 0.5-1.0 ug/mL Optimal dilution for a specific application should be determined by user |