

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Ep-CAM / CD326 (Extracellular Domain) (Epithelial Marker)(MOC-31), 1mg/mL, Clone: [MOC-31], Mouse, Monoclonal BOT-BNUM0380-50

Article Name	Ep-CAM / CD326 (Extracellular Domain) (Epithelial Marker)(MOC-31), 1mg/mL, Clone: [MOC-31], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUM0380-50
Supplier Catalog Number	BNUM0380-50
Alternative Catalog Number	BOT-BNUM0380-50-50UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC, WB
Species Reactivity	Human
Immunogen	Neuraminidase treated GLS-1 human small cell lung carcinoma cells
Product Description	The epitope of this antibody is located in the first EGF-like repeat domain (EGF1) between amino acids 27-59 of Ep-CAM. EGP40 is a 40-43 kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cel...
Clonality	Monoclonal
Concentration	1 mg/mL
Clone Designation	[MOC-31]
Molecular Weight	40-43 kDa

UniProt	P16422
Buffer	PBS, no BSA, no azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistochemistry (formalin-fixed): 1-2 ug/mL for 30 minutes at RT Western blot: 1-2 ug/mL Does not react with rat Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user