

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Thomsen-Friedenreich Antigen / CD176 (Pan Carcinoma Marker)(A63-C/A9), 1mg/mL, Clone: [A63-C/A9], Mouse, Monoclonal BOT-BNUM0938-50

Article Name	Thomsen-Friedenreich Antigen / CD176 (Pan Carcinoma Marker)(A63-C/A9), 1mg/mL, Clone: [A63-C/A9], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUM0938-50
Supplier Catalog Number	BNUM0938-50
Alternative Catalog Number	BOT-BNUM0938-50-50UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human, Mouse, Rat
Immunogen	Neuraminidase-treated human red blood cells
Product Description	This antibody recognizes a disaccharide epitope, Gal 1-3GalNAc, of Thomsen-Friedenreich (TF) antigen. It is specific for both anomeric forms of the disaccharide (TF and TF , including related structures on the glycolipid) and shows no cross-reactivit...
Clonality	Monoclonal
Concentration	1 mg/mL
Clone Designation	[A63-C/A9]
Molecular Weight	Multiple

UniProt	Not Applicable
Buffer	PBS, no BSA, no azide
Source	Animal
Application Notes	Immunohistochemistry (formalin-fixed): 1-2 ug/mL for 30 minutes at 37°C Immunohistochemistry: frozen: 0.5-1.0 ug/mL for 30 minutes at RT Immunofluorescence: 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user