

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Cytokeratin, LMW(KRTL/1077), 1mg/mL, Clone: [KRTL/1077], Mouse, Monoclonal BOT-BNUM1077-50

Article Name	Cytokeratin, LMW(KRTL/1077), 1mg/mL, Clone: [KRTL/1077], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUM1077-50
Supplier Catalog Number	BNUM1077-50
Alternative Catalog Number	BOT-BNUM1077-50-50UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human, Rat
Immunogen	Recombinant human KRT77 protein
Product Description	This MAb recognizes the 56.5 kDa (CK10), 50 kDa (CK14), 50 kDa (CK15), 48 kDa (CK16), 40 kDa (CK19) keratins of the acidic (Type I or LMW) subfamily. Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0) sub...
Clonality	Monoclonal
Concentration	1 mg/mL
Clone Designation	[KRTL/1077]
Molecular Weight	56.5 kDa (CK10), 50 kDa (CK14), 50 kDa (CK15), 48 kDa (CK16), 40 kDa (CK19)

UniProt	Q7Z794
Buffer	PBS, no BSA, no azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Predicted to show broad species reactivity Optimal dilution for a specific application should be determined by user