

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CDw17(H018.3G-6.F5), 1mg/mL, IgM, Clone: [H018.3G-6.F5], Mouse, Monoclonal BOT-BNUM1092-50

Article Name	CDw17(H018.3G-6.F5), 1mg/mL, IgM, Clone: [H018.3G-6.F5], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUM1092-50
Supplier Catalog Number	BNUM1092-50
Alternative Catalog Number	BOT-BNUM1092-50-50UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Species Reactivity	Human
Immunogen	?-2 Microglobulin associated proteins from a detergent lysate of human PBL
Product Description	CDw17 / CD17 is an intermediate glycosphingolipid from the metabolism of higher gangliosides that localizes to sphingolipid-sterol rafts. CD17 is detectable in monocytes, granulocytes, basophils, platelets, a subset of peripheral B cells (CD19) and ...
Clonality	Monoclonal
Concentration	1 mg/mL
Clone Designation	[H018.3G-6.F5]
Molecular Weight	Not Known
Isotype	IgM

UniProt	Not Known
Buffer	PBS, no BSA, no azide
Source	Animal
Application Notes	For coating for ELISA, order Ab without BSA Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry