

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD45RO(UCHL1 + T200/797), 1mg/mL, Clone: [UCHL-1 + T200/797], Mouse, Monoclonal BOT-BNUM1209-50

| | |
|----------------------------|---|
| Article Name | CD45RO(UCHL1 + T200/797), 1mg/mL, Clone: [UCHL-1 + T200/797], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNUM1209-50 |
| Supplier Catalog Number | BNUM1209-50 |
| Alternative Catalog Number | BOT-BNUM1209-50-50UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC, WB |
| Species Reactivity | Human |
| Immunogen | Cultured human T-cells from an IL-2-dependent T-cell line (CA1) |
| Product Description | Recognizes a 180-185 kDa protein, identified as isoform of leukocyte common antigen (CD45RO). This antibody reacts with mature activated T-cells, most thymocytes, and a sub-population of resting T-cells within both CD4 and CD8 subsets. It shows no re... |
| Clonality | Monoclonal |
| Concentration | 1 mg/mL |
| Clone Designation | [UCHL-1 + T200/797] |
| Molecular Weight | 180-185 kDa |

| | |
|-------------------|---|
| UniProt | P08575 |
| Buffer | PBS, no BSA, no azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |