

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

VEGI (Vascular Endothelial Growth Inhibitor) (VEGI/1283), 1mg/mL, Clone: [VEGI/1283], Mouse, Monoclonal BOT-BNUM1283-50

| | |
|----------------------------|---|
| Article Name | VEGI (Vascular Endothelial Growth Inhibitor) (VEGI/1283), 1mg/mL, Clone: [VEGI/1283], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNUM1283-50 |
| Supplier Catalog Number | BNUM1283-50 |
| Alternative Catalog Number | BOT-BNUM1283-50-50UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant human full-length VEGI fragment |
| Product Description | VEGI is an anti-angiogenic cytokine that belongs to tumor necrosis factor superfamily, member 15 (TNFSF15). This protein is abundantly expressed in endothelial cells, but is not expressed in either B or T cells. The expression of this protein is indu... |
| Clonality | Monoclonal |
| Concentration | 1 mg/mL |
| Clone Designation | [VEGI/1283] |
| Molecular Weight | 22 kDa |

| | |
|-------------------|---|
| UniProt | O95150 |
| Buffer | PBS, no BSA, no azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Flow cytometry: 0.5-1 ug/million cells in 0.1mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Western blotting 1-2 ug/mL Optimal dilution for a specific application should be determined by user |