

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD1a / HTA1 (Mature Langerhans Cells Marker) (C1A/1506R), 1mg/mL, Clone: [C1A/1506R], Rabbit, Monoclonal BOT-BNUM1506-50

Article Name	CD1a / HTA1 (Mature Langerhans Cells Marker) (C1A/1506R), 1mg/mL, Clone: [C1A/1506R], Rabbit, Monoclonal
Biozol Catalog Number	BOT-BNUM1506-50
Supplier Catalog Number	BNUM1506-50
Alternative Catalog Number	BOT-BNUM1506-50-50UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant human full-length CD1a protein
Product Description	At least five CD1 genes (CD1a, b, c, d, and e) are identified. CD1 proteins have been demonstrated to restrict T cell response to non-peptide lipid and glycolipid antigens and play a role in non-classical antigen presentation. CD1a is a non-polymorph...
Clonality	Monoclonal
Concentration	1 mg/mL
Clone Designation	[C1A/1506R]
Molecular Weight	49 kDa

UniProt	P06126
Buffer	PBS, no BSA, no azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user