

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

TCL1 (T-Cell Marker) (TCL1/2078), 1mg/mL, Clone: [TCL1/2078], Mouse, Monoclonal BOT-BNUM2078-50

Article Name	TCL1 (T-Cell Marker) (TCL1/2078), 1mg/mL, Clone: [TCL1/2078], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUM2078-50
Supplier Catalog Number	BNUM2078-50
Alternative Catalog Number	BOT-BNUM2078-50-50UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant human TCL1 protein fragment (around aa 2-109) (exact sequence is proprietary)
Product Description	T-cell leukemia/Lymphoma Protein 1A (TCL-1A), also known as p14TCL1 is a product of the TCL1 gene that is involved in T-cell prolymphocytic leukemia (TPLL). T-PLL is a rare form of mature T-cell leukemia, which is consistently associated with chromos...
Clonality	Monoclonal
Concentration	1 mg/mL
Clone Designation	[TCL1/2078]
Molecular Weight	14 kDa

UniProt	P56279
Buffer	PBS, no BSA, no azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user