

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Bcl-2 (Apoptosis & Follicular Lymphoma Marker) (BCL2/2210R), 1mg/mL, Clone: [BCL2/2210R], Rabbit, Monoclonal BOT-BNUM2210-50

| | |
|----------------------------|---|
| Article Name | Bcl-2 (Apoptosis & Follicular Lymphoma Marker) (BCL2/2210R), 1mg/mL, Clone: [BCL2/2210R], Rabbit, Monoclonal |
| Biozol Catalog Number | BOT-BNUM2210-50 |
| Supplier Catalog Number | BNUM2210-50 |
| Alternative Catalog Number | BOT-BNUM2210-50-50UL |
| Manufacturer | Biotium |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC, WB |
| Species Reactivity | Human |
| Immunogen | Recombinant full-length human bcl-2 protein |
| Product Description | This antibody recognizes a protein of 25-26 kDa, identified as Bcl-2. This antibody shows no cross-reaction with Bcl-x or Bax proteins. Bcl-2 suppresses apoptosis in a variety of cell systems including factor-dependent lymphohematopoietic and neural ... |
| Clonality | Monoclonal |
| Concentration | 1 mg/mL |
| Clone Designation | [BCL2/2210R] |
| Molecular Weight | 25-26 kDa |

| | |
|-------------------|--|
| UniProt | P10415 |
| Buffer | PBS, no BSA, no azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris with 1 mM EDTA Buffer pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |