

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Tenascin C (Stromal Marker For Epithelial Malignancy) (TNC/2981R), 1mg/mL, Clone: [TNC/2981R], Rabbit, Monoclonal BOT-BNUM2981-50

Article Name	Tenascin C (Stromal Marker For Epithelial Malignancy) (TNC/2981R), 1mg/mL, Clone: [TNC/2981R], Rabbit, Monoclonal
Biozol Catalog Number	BOT-BNUM2981-50
Supplier Catalog Number	BNUM2981-50
Alternative Catalog Number	BOT-BNUM2981-50-50UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant full-length human TNC protein
Product Description	In Western blotting, this antibody reacts with two bands of ~MW of 210 kDa and 300 kDa, identified as two isoforms of Tenascin C. Specificity of this MAb is validated by sequential immunoprecipitation with a PAb against Tenascin C. Tenascin C is a mu...
Clonality	Monoclonal
Concentration	1 mg/mL
Clone Designation	[TNC/2981R]
Molecular Weight	210 kDa and 300 kDa

UniProt	P24821
Buffer	PBS, no BSA, no azide
Source	Animal
Application Notes	<p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Does not react with rat Immunohistology (formalin): 2-4 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user</p>