

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **BCL10 (MALT-Lymphoma Marker) (BL10/2988R), 1mg/mL, Clone: [BL10/2988R], Rabbit, Monoclonal BOT-BNUM2988-50**

Article Name	BCL10 (MALT-Lymphoma Marker) (BL10/2988R), 1mg/mL, Clone: [BL10/2988R], Rabbit, Monoclonal
Biozol Catalog Number	BOT-BNUM2988-50
Supplier Catalog Number	BNUM2988-50
Alternative Catalog Number	BOT-BNUM2988-50-50UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Application	FC, IF, IHC
Species Reactivity	Human
Immunogen	Recombinant human BCL10 protein fragment (around aa122-168) (exact sequence is proprietary)
Product Description	B-cell lymphoma/leukemia 10 (BCL10) is a signaling protein that promotes apoptosis, pro-caspase-9 maturation, and activation of NF-kappa-B via NIK and IKK. BCL10 contains an N-terminal caspase recruitment domain (CARD), which is found in a number of ...
Clonality	Monoclonal
Concentration	1 mg/mL
Clone Designation	[BL10/2988R]
Molecular Weight	33 kDa

UniProt	O95999
Buffer	PBS, no BSA, no azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user