

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Recombinant Human Tumor Necrosis Factor-alpha (rHuTNF-alpha) BWT-PR1125

| Article Name               | Recombinant Human Tumor Necrosis Factor-alpha (rHuTNF-alpha)   |
|----------------------------|--|
| Biozol Catalog Number      | BWT-PR1125   |
| Supplier Catalog Number    | PR1125   |
| Alternative Catalog Number | BWT-PR1125-10UG,BWT-PR1125-50UG,BWT-PR1125-1.0MG   |
| Manufacturer               | Bioworld Technology  |
| Category                   | Biochemikalien   |
| Product Description        | Tumor necrosis factor alpha (TNF-alpha), also called cachectin, is produced by neutrophils, activated lymphocytes, macrophages, NK cells, LAK cells, astrocytes endothelial cells, smooth muscle cells and some transformed cells. TNF-alpha occurs as a s |
| Molecular Weight           | Approximately 17.5 kDa. The recombinant protein preparation is a mixture of a 158 amino acid residue form containing an N-terminal methionine and a 157 amino acid residue form with the sequence of mature human TNF-alpha.                               |
| Source                     | Escherichia coli.  |
| Purity                     | >95% by SDS-PAGE and HPLC analyses.  |
| Form                       | Sterile Filtered White lyophilized (freeze-dried) powder.  |
| Sequence                   | VRSSSRTPSD KPVAHVVANP QAEGQLQWLN RRANALLANG VELRDNQLVVPSEGLYLIYS QVLFKGQGCP STHVLLTHTI SRIAVSYQTK VNLLSAIKSP CQRETPEGAE AKPWYEPIYL GGVFQLEKGD RLSAEINRPD YLDFAESGQV YFGIIAL  |
| Formula                    | Lyophilized from a 0.2mm filtered concentrated solution in PBS, pH 7.0.  |

## **Application Notes**

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportio