

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Alexa Fluor 594-conjugated AffiniPure Donkey Anti-Rabbit IgG (H+L) (min X Bov,Ck,Gt,GP,Sy Hms,Hrs,Hu,Ms,Rat,Shp Sr Prot) JIM-711-585-152**

|                            |   |
|----------------------------|---|
| Article Name               | Alexa Fluor 594-conjugated AffiniPure Donkey Anti-Rabbit IgG (H+L)<br>(min X Bov,Ck,Gt,GP,Sy Hms,Hrs,Hu,Ms,Rat,Shp Sr Prot)                                     |
| Biozol Catalog Number      | JIM-711-585-152   |
| Supplier Catalog Number    | 711-585-152   |
| Alternative Catalog Number | JIM-711-585-152   |
| Manufacturer               | Jackson ImmunoResearch  |
| Host                       | Donkey  |
| Category                   | Antikörper  |
| Species Reactivity         | Rabbit  |
| Conjugation                | Alexa Fluor 594   |
| Format                     | IgG   |
| Target Specificity         | IgG (H+L)   |
| Cross-Adsorption (MinX)    | Bovine,Gallus,Goat,Guinea pig,Hamster<br>(syrian),Equine,Human,Mouse,Rat,Sheep  |
| Clonality                  | Polyclonal  |
| Buffer                     | Buffer: 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6. Stabilizer: 15<br>mg/ml Bovine Serum Albumin (IgG-Free, Protease-Free).<br>Preservative: 0.05% Sodium Azide |
| Purity                     | AffiniPure  |

|                    |  |
|--------------------|--|
| Form               | Freeze-dried solid   |
| Storage            | Storage and Rehydration: Store freeze-dried solid at 2-8°C. Rehydrate with the indicated volume of dH <sub>2</sub> O and centrifuge if not clear. Prepare working dilution on day of use. Product is stable for about 6 weeks at 2-8°C as an undiluted liquid. Extended... |
| Antibody Type      | Secondary Antibody   |
| Application Dilute | 1:100 - 1:800 for most applications Dilution factors are presented in the form of a range because the optimal dilution is a function of many factors, such as antigen density, permeability, etc. The actual dilution used must be determined empirically.                 |