

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Complement Factor D Antibody / CFD, Rabbit, Polyclonal NSJ-F54234-0.4ML

Article Name Complement Factor D Antibody / CFD, Rabbit, Polyclonal Biozol Catalog Number NSJ-F54234-0.4ML Supplier Catalog Number F54234-0.4ML Alternative Catalog Number NSJ-F54234-0.4ML Manufacturer NSJ Bioreagents Host Rabbit Category Antikörper Application IHC-P, WB Species Reactivity Human Immunogen A portion of amino acids 68-99 from the human protein were used at the immunogen for the Complement Factor D antibody. The protein encoded by this gene is a member of the trypsin family of peptidases. The encoded protein is a component of the alternative complement pathway best known for its role in humoral suppressio
Supplier Catalog Number F54234-0.4ML Alternative Catalog Number NSJ-F54234-0.4ML Manufacturer NSJ Bioreagents Host Rabbit Category Antikörper Application IHC-P, WB Species Reactivity Human Immunogen A portion of amino acids 68-99 from the human protein were used at the immunogen for the Complement Factor D antibody. The protein encoded by this gene is a member of the trypsin family of peptidases. The encoded protein is a component of the alternative
Alternative Catalog Number NSJ-F54234-0.4ML Manufacturer NSJ Bioreagents Host Rabbit Category Antikörper Application IHC-P, WB Species Reactivity Human A portion of amino acids 68-99 from the human protein were used at the immunogen for the Complement Factor D antibody. The protein encoded by this gene is a member of the trypsin family of peptidases. The encoded protein is a component of the alternative
Manufacturer NSJ Bioreagents Host Rabbit Category Antikörper Application IHC-P, WB Species Reactivity Human A portion of amino acids 68-99 from the human protein were used a the immunogen for the Complement Factor D antibody. The protein encoded by this gene is a member of the trypsin family of peptidases. The encoded protein is a component of the alternative
Host Rabbit Category Antikörper Application IHC-P, WB Species Reactivity Human Immunogen A portion of amino acids 68-99 from the human protein were used at the immunogen for the Complement Factor D antibody. The protein encoded by this gene is a member of the trypsin family of peptidases. The encoded protein is a component of the alternative
Category Application IHC-P, WB Species Reactivity Human A portion of amino acids 68-99 from the human protein were used a the immunogen for the Complement Factor D antibody. The protein encoded by this gene is a member of the trypsin family of peptidases. The encoded protein is a component of the alternative.
Application IHC-P, WB Species Reactivity Human A portion of amino acids 68-99 from the human protein were used at the immunogen for the Complement Factor D antibody. The protein encoded by this gene is a member of the trypsin family of peptidases. The encoded protein is a component of the alternative
Species Reactivity Human A portion of amino acids 68-99 from the human protein were used a the immunogen for the Complement Factor D antibody. The protein encoded by this gene is a member of the trypsin family of peptidases. The encoded protein is a component of the alternative
A portion of amino acids 68-99 from the human protein were used at the immunogen for the Complement Factor D antibody. The protein encoded by this gene is a member of the trypsin family of peptidases. The encoded protein is a component of the alternative
The protein encoded by this gene is a member of the trypsin family of peptidases. The encoded protein is a component of the alternative
Product Description of peptidases. The encoded protein is a component of the alternative
of infectious agents. This protein is also a ser
Clonality Polyclonal
UniProt P00746
Purity Antigen affinity purified
Form In 1X PBS, pH 7.4, with 0.09% sodium azide
Antibody Type Primary Antibody

Application Dilute	Western blot: 1:1000-1:2000,Immunohistochemistry: 1:50-1:100
Application Notes	The stated application concentrations are suggested starting points. Titration of the Complement Factor D antibody may be required due to differences in protocols and secondary/substrate sensitivity.