

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Rat Coagulation Factor II protein, His tag, Unconjugated GTX00058-PRO

Article Name	Rat Coagulation Factor II protein, His tag, Unconjugated
Biozol Catalog Number	GTX00058-PRO
Supplier Catalog Number	GTX00058-pro
Alternative Catalog Number	GTX00058-PRO-10
Manufacturer	GeneTex
Category	Proteine/Peptide
Application	FA
Species Reactivity	Rat
Conjugation	Unconjugated
NCBI	<a href="#">29251</a>
UniProt	<a href="#">P18292</a>
Buffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Form	Lyophilized powder
Sequence	N-terminal His-Tag, Ser201~Arg323

Application Notes

Coagulation Factor II (F2) also commonly called pro-thrombin is a coagulation protein in the blood stream that has many effects in the coagulation cascade. It is a serine protease that converts fibrinogen to fibrin and activates factors V, VII, VIII, XIII, and, in complex with thrombomodulin, protein C. Besides, Protein C Inhibitor (PCI) has been identified as an interactor of F2, thus a binding ELISA assay was conducted to detect the interaction of recombinant rat F2 and recombinant rat PCI. Briefly, F2 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100  $\mu$ l were then transferred to PCI-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-F2 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50  $\mu$ l stop solution to the wells and read at 450nm immediately. The binding activity of F2 and PCI was in a dose dependent manner.