

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Human HMGB1 protein, His tag, Unconjugated GTX00153-PRO

Article Name	Human HMGB1 protein, His tag, Unconjugated
Biozol Catalog Number	GTX00153-PRO
Supplier Catalog Number	GTX00153-pro
Alternative Catalog Number	GTX00153-PRO-10
Manufacturer	GeneTex
Category	Proteine/Peptide
Application	FA
Species Reactivity	Human
Conjugation	Unconjugated
NCBI	3146
UniProt	P09429
Buffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Form	Lyophilized powder
Sequence	N-terminal His-Tag, Pro9~Arg163 (NP_001300821.1)

Application Notes

High Mobility Group Protein 1 (HMG1) is among the most important chromatin proteins. In the nucleus HMG1 interacts with nucleosomes, transcription factors, and histones. This nuclear protein organizes the DNA and regulates transcription. Besides, Stromal Cell Derived Factor 1 (SDF1) has been identified as an interactor of HMG1, thus a binding ELISA assay was conducted to detect the interaction of recombinant human HMG1 and recombinant human SDF1. Briefly, HMG1 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 μ l were then transferred to SDF1-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-HMG1 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 μ l stop solution to the wells and read at 450nm immediately. The binding activity of HMG1 and SDF1 was in a dose dependent manner.