

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Human CYP3A4 protein, His tag, Unconjugated GTX00198-PRO

Article Name	Human CYP3A4 protein, His tag, Unconjugated
Biozol Catalog Number	GTX00198-PRO
Supplier Catalog Number	GTX00198-pro
Alternative Catalog Number	GTX00198-PRO-10
Manufacturer	GeneTex
Category	Proteine/Peptide
Application	FA
Species Reactivity	Human
Conjugation	Unconjugated
NCBI	1576
UniProt	P08684
Buffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Form	Lyophilized powder
Sequence	N-terminal His-Tag, Pro344~Asp497@ (NP_001189784.1)

Application Notes

Cytochrome P450 3A4 (CYP3A4) is an important enzyme in the body, mainly found in the liver and in the intestine. It oxidizes small foreign organic molecules (xenobiotics), such as toxins or drugs, so that they can be removed from the body. CYP3A4 is a member of the cytochrome P450 family of oxidizing enzymes and the CYP3A4 is the most common and the most versatile one in drug metabolism. Besides, Heat Shock 70kDa Protein 8 (HSPA8) has been identified as an interactor of CYP3A4, thus a binding ELISA assay was conducted to detect the interaction of recombinant human CYP3A4 and recombinant human HSPA8. Briefly, CYP3A4 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 μ l were then transferred to HSPA8-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-CYP3A4 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of CYP3A4 and HSPA8 was in a dose dependent manner.