

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Human CHI3L1 protein, His tag (active), Unconjugated GTX00219-PRO

| Article Name | Human CHI3L1 protein, His tag (active), Unconjugated |
|----------------------------|---|
| Biozol Catalog Number | GTX00219-PRO |
| Supplier Catalog Number | GTX00219-pro |
| Alternative Catalog Number | GTX00219-PRO-10 |
| Manufacturer | GeneTex |
| Category | Proteine/Peptide |
| Application | FA |
| Species Reactivity | Human |
| Conjugation | Unconjugated |
| NCBI | 1116 |
| UniProt | P36222 |
| Buffer | Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300. |
| Expression System | E. coli |
| Form | Lyophilized powder |
| Sequence | N-terminal His-Tag, Asn112~Thr237+Ile299~Lys377 (NP_001267.2) |

Application Notes

Cell proliferation of Raji cells after stimulated with GP39. Cell proliferation of Raji cells after stimulated with GP39. Glycoprotein 39, Cartilage (GP39) also known as Chitinase-3-like protein 1 (CHI3L1) is a secreted glycoprotein that is approximately 40kDa in size that in humans is encoded by the CHI3L1 gene. GP39 plays a role in cancer cell proliferation, survival, invasiveness and in the regulation of cellmatrix interactions. To test the effect of GP39 on cell proliferation, Raji cells were seeded into triplicate wells of 96-well plates at a density of 5000 cells/well with 1% serum standard RPMI-1640 which contains various concentrations of recombinant human GP39. After incubated for 5 days, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10 µl of CCK-8 solution was added to each well of the plate, then the absorbance at 450nm was measured using a microplate reader after incubating the plate for 1-4 hours at 37C. Proliferation of Raji cells after incubation with GP39 for 5 days observed by inverted microscope.