

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Human MX1 protein, His tag, Unconjugated GTX00244-PRO

Article Name	Human MX1 protein, His tag, Unconjugated
Biozol Catalog Number	GTX00244-PRO
Supplier Catalog Number	GTX00244-pro
Alternative Catalog Number	GTX00244-PRO-10
Manufacturer	GeneTex
Category	Proteine/Peptide
Application	FA
Species Reactivity	Human
Conjugation	Unconjugated
NCBI	4599
UniProt	P20591
Buffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Form	Lyophilized powder
Sequence	N-terminal His-Tag, Ser80~Leu342 (NP_001138397.1)

Application Notes

Interferon-induced GTP-binding protein Mx1 is a protein that in humans is encoded by the MX1 gene. In mouse, the interferon-inducible Mx protein is responsible for a specific antiviral state against influenza virus infection. The protein encoded by this gene is similar to the mouse protein as determined by its antigenic relatedness, induction conditions, physicochemical properties, and amino acid analysis. This cytoplasmic protein is a member of both the dynamin family and the family of large GTPases. Besides, Tubulin Beta (TUBb) has been identified as an interactor of MX1, thus a binding ELISA assay was conducted to detect the interaction of recombinant human MX1 and recombinant human TUBb. Briefly, MX1 was diluted serially in PBS with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl were then transferred to TUBb-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-MX1 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of MX1 and TUBb was in a dose dependent manner.