

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Human IGF1 protein, His tag (active), Unconjugated GTX00250-PRO

Article Name	Human IGF1 protein, His tag (active), Unconjugated
Biozol Catalog Number	GTX00250-PRO
Supplier Catalog Number	GTX00250-pro
Alternative Catalog Number	GTX00250-PRO-10
Manufacturer	GeneTex
Category	Proteine/Peptide
Application	FA
Species Reactivity	Human
Conjugation	Unconjugated
NCBI	3479
UniProt	P05019
Buffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Form	Lyophilized powder
Sequence	N-terminal His-Tag, Gly49~Ala118 (NP_000609.1)

Application Notes

insulin-like growth factor 1 (IGF1), also called somatomedin C is a hormone similar in molecular structure to insulin. It plays an important role in childhood growth and continues to have anabolic effects in adults. A synthetic analog of IGF1, mecasermin, is used for the treatment of growth failure. IGF1 consists of 70 amino acids in a single chain with three intramolecular disulfide bridges. IGF1 has a molecular weight of 7, 649 Daltons. Besides, insulin Like Growth Factor Binding Protein 4 (IGFBP4) has been identified as an interactor of IGF1, thus a binding ELISA assay was conducted to detect the interaction of recombinant human IGF1 and recombinant human IGFBP4. Briefly, IGF1 were diluted serially in PBS, with 0. 01% BSA (pH 7. 4). Duplicate samples of 100 μ l were then transferred to IGFBP4-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-VDR pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 μ l stop solution to the wells and read at 450nm immediately. The binding activity of IGF1 and IGFBP4 was in a dose dependent manner. To measure the effect of IGF1 on cell proliferation, breast cancer MCF-7 cells were seeded into triplicate wells of 96-well plates at a density of 5000 cells/well and allowed to attach, replaced with serum-free overnight, then the medium was replaced with 1% serum standard DMEM prior to the addition of various concentrations of recombinant human IGF1. After incubated for 96h, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10 μ l of CCK-8 solution was added to each well of the plate, then the absorbance at 450nm was measured using a microplate reader after incubating the plate for 1-4 hours at 37C. Proliferation of MCF-7 cells after incubation with IGF1 for 72h observed by inverted microscop. Cell viability was assessed by CCK-8 assay after incubation with recombinant IGF1 for 72h, and IGF1 significantly increased cell viability of MCF-7 cells.