

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Mouse TNF alpha protein, His tag (active), Unconjugated GTX00288-PRO

| Article Name | Mouse TNF alpha protein, His tag (active), Unconjugated |
|----------------------------|---|
| Biozol Catalog Number | GTX00288-PRO |
| Supplier Catalog Number | GTX00288-pro |
| Alternative Catalog Number | GTX00288-PRO-10 |
| Manufacturer | GeneTex |
| Category | Proteine/Peptide |
| Application | FA |
| Species Reactivity | Mouse |
| Conjugation | Unconjugated |
| NCBI | 21926 |
| UniProt | P06804 |
| Buffer | Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300. |
| Expression System | HEK293 cells |
| Form | Lyophilized powder |
| Sequence | N-terminal His-Tag, Gly57~Leu235 (NP_001265530.1) |

Application Notes

TNF-a (Tumor necrosis factor), being an endogenous pyrogen, is able to induce fever, necrosis, inflammation and to inhibit tumorigenesis. As reported, TNF-a could inhibit the proliferation and induce necrosis of A549 cells, and the concentration of IL-1beta in cell supernatant will increase after stimulation. Therefore, A549 cells were incubated in DMEM with TNFa (1 ng/ml, 10 ng/ml) for 2h, 4h, 8h, 24h, 48h, then cells were observed by inverted microscope. Cell necrosis after incubation with TNF-a (10 ng/ml) for 72h.