

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Mouse LCAT protein, His tag, Unconjugated GTX00300-PRO

Article Name	Mouse LCAT protein, His tag, Unconjugated
Biozol Catalog Number	GTX00300-PRO
Supplier Catalog Number	GTX00300-pro
Alternative Catalog Number	GTX00300-PRO-10
Manufacturer	GeneTex
Category	Proteine/Peptide
Application	FA
Species Reactivity	Mouse
Conjugation	Unconjugated
NCBI	16816
UniProt	P16301
Buffer	Reconstitute with 10mM PBS (pH7.4) to 0.1-1.0mg/ml. Do not vortex. Lyophilized from PBS (pH7.4), 0.01% SKL, 1mM DTT, 5% Trehalose, ProClin 300.
Expression System	E. coli
Form	Lyophilized powder
Sequence	N-terminal His-Tag, Glu41~His210 (NP_032516.2)

**Application Notes** 

LCAT (Phosphatidylcholine-sterol acyltransferase) is an enzyme in the extracellular metabolism of plasma lipoproteins, which converts cholesterol and phosphatidylcholines (lecithins) to cholesteryl esters and lysophosphatidylcholines. It is reported that APOA1 (Apolipoprotein A-I) acts as a cofactor for the LCAT. Thus, a binding ELISA assay was conducted to detect the association of LCAT with APOA1. Briefly, recombinant rat LCAT were diluted serially in PBS with 0.01% BSA (pH 7.4). Duplicate samples of 100 μl were then transferred to APOA1-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-LCAT pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50  $\mu$ l stop solution to the wells and read at 450nm immediately. The binding activity of LCAT with APOA1 was in a dose dependent manner.