

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Mouse LIF protein, His tag, Unconjugated GTX00320-PRO

Article Name	Mouse LIF protein, His tag, Unconjugated
Biozol Catalog Number	GTX00320-PRO
Supplier Catalog Number	GTX00320-pro
Alternative Catalog Number	GTX00320-PRO-10
Manufacturer	GeneTex
Category	Proteine/Peptide
Application	FA
Species Reactivity	Mouse
Conjugation	Unconjugated
NCBI	16878
UniProt	P09056
Buffer	Reconstitute with 10mM PBS (pH7.4) to 0.1-1.0mg/ml. Do not vortex. Lyophilized from PBS (pH7.4), 0.01% SKL, 1mM DTT, 5% Trehalose, ProClin 300.
Expression System	E. coli
Form	Lyophilized powder
Sequence	N-terminal His-Tag, Pro25~Phe203 (NP_001034626.1)

Application Notes	<p>Leukemia inhibitory factor (LIF), is an interleukin 6 class cytokine that affects cell growth by inhibiting differentiation. LIF as a cytokine also has another function including: the growth promotion and cell differentiation of different types of target cells, influence on bone metabolism, cachexia, neural development, embryogenesis and inflammation. Besides, Colony Stimulating Factor Receptor, Granulocyte (GCSFR) has been identified as an interactor of LIF, thus a binding ELISA assay was conducted to detect the interaction of recombinant mouse LIF and recombinant mouse GCSFR. Briefly, LIF were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl were then transferred to GCSFR-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-LIF pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of LIF and GCSFR was in a dose dependent manner.</p>
-------------------	---