



Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Mouse iNOS protein, His tag, Unconjugated GTX00325-PRO

Article Name	Mouse iNOS protein, His tag, Unconjugated
Biozol Catalog Number	GTX00325-PRO
Supplier Catalog Number	GTX00325-pro
Alternative Catalog Number	GTX00325-PRO-10
Manufacturer	GeneTex
Category	Proteine/Peptide
Application	FA
Species Reactivity	Mouse
Conjugation	Unconjugated
NCBI	18126
UniProt	P29477
Buffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Form	Lyophilized powder
Sequence	N-terminal His-Tag, Asp43~Thr213 (NP_001300850.1)

Application Notes

Nitric oxide synthase 2, inducible (NOS2) is a member of Nitric oxide synthases (NOSs) family. Nitric oxide synthases (NOSs) are a family of enzymes catalyzing the production of nitric oxide (NO) from Larginine. NO is an important cellular signaling molecule. It helps modulate vascular tone, insulin secretion, airway tone, and peristalsis, and is involved in angiogenesis and neural development. Besides, Ubiquitin Carboxyl Terminal Hydrolase L5 (UCHL5) has been identified as an interactor of NOS2, thus a binding ELISA assay was conducted to detect the interaction of recombinant mouse NOS2 and recombinant mouse UCHL5. Briefly, NOS2 was diluted serially in PBS with 0.01% BSA (pH 7.4). Duplicate samples of 100 μl were then transferred to UCHL5-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-NOS2 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 μl stop solution to the wells and read at 450nm immediately. The binding activity of NOS2 and UCHL5 was in a dose dependent manner.