

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rat CXCL2 / GRO beta protein, His and GST tag, Unconjugated GTX00351-PRO

| Article Name | Rat CXCL2 / GRO beta protein, His and GST tag, Unconjugated |
|----------------------------|---|
| Biozol Catalog Number | GTX00351-PRO |
| Supplier Catalog Number | GTX00351-pro |
| Alternative Catalog Number | GTX00351-PRO-10 |
| Manufacturer | GeneTex |
| Category | Proteine/Peptide |
| Application | FA |
| Species Reactivity | Rat |
| Conjugation | Unconjugated |
| NCBI | 114105 |
| UniProt | P30348 |
| Buffer | Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300. |
| Expression System | E. coli |
| Form | Lyophilized powder |
| Sequence | N-terminal His and GST-Tag, Ser32~Asn100 (NP_446099.1) |

Application Notes

Growth Regulated Oncogene Beta (GROb) is a small cytokine belonging to the CXC chemokine family which produced by activated monocytes and neutrophils and expressed at sites of inflammation. Hematoregulatory chemokine, which, in vitro, suppresses hematopoietic progenitor cell proliferation. Besides, Dipeptidyl Peptidase IV (DPP4) has been identified as an interactor of GROb, thus a binding ELISA assay was conducted to detect the interaction of recombinant rat GROb and recombinant rat DPP4. Briefly, GROb were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl were then transferred to DPP4-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-GROb pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 μl stop solution to the wells and read at 450nm immediately. The binding activity of of GROb and DPP4 was in a dose dependent manner.