

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rat Plasminogen protein, His tag, Unconjugated GTX00354-PRO

Article Name	Rat Plasminogen protein, His tag, Unconjugated
Biozol Catalog Number	GTX00354-PRO
Supplier Catalog Number	GTX00354-pro
Alternative Catalog Number	GTX00354-PRO-10
Manufacturer	GeneTex
Category	Proteine/Peptide
Application	FA
Species Reactivity	Rat
Conjugation	Unconjugated
NCBI	85253
UniProt	Q01177
Buffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Form	Lyophilized powder
Sequence	N-terminal His-Tag, Glu191~Arg433 (NP_445943.1)

Application Notes

Plasminogen (Plg) can be converted into active plasmin by tissue plasminogen activator (tPA), urokinase plasminogen activator (uPA), kallikrein, and factor XII (Hageman factor). Plasmin can dissolve fibrin blood clots, act on many other processes such as embryonic development, tissue remodeling, inflammation and tumor invasion. Plasmin also activates collagenases, weakens the walls of the Graafian follicle, cleaves fibrin, fibronectin, thrombospondin, laminin, and von Willebrand factor. Besides, Actin Beta (ACTb) has been identified as an interactor of Plg, thus a binding ELISA assay was conducted to detect the interaction of recombinant rat Plg and recombinant rat ACTb. Briefly, Plg were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl were then transferred to ACTb-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-Plg pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of Plg and ACTb was in a dose dependent manner.