

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **TLE1 (Synovial Sarcoma Marker) (TLE1/2062), CF740 conjugate, 0.1mg/mL, Clone: [TLE1/2062], CF 740, Mouse, Monoclonal BOT-BNC742062-500**

|                            |   |
|----------------------------|---|
| Article Name               | TLE1 (Synovial Sarcoma Marker) (TLE1/2062), CF740 conjugate, 0.1mg/mL, Clone: [TLE1/2062], CF 740, Mouse, Monoclonal  |
| Biozol Catalog Number      | BOT-BNC742062-500   |
| Supplier Catalog Number    | BNC742062-500   |
| Alternative Catalog Number | BOT-BNC742062-500-500UL   |
| Manufacturer               | Biotium   |
| Host                       | Mouse   |
| Category                   | Antikörper  |
| Application                | IHC, WB   |
| Species Reactivity         | Human   |
| Immunogen                  | Recombinant human TLE1 fragment (aa 175-338) (exact sequence is proprietary)  |
| Conjugation                | CF 740  |
| Product Description        | Key players in the Notch pathway are the TLE genes, which are human homologs of the Drosophila groucho gene. Groucho is a transcriptional repressor that plays a key role in neurogenesis, segmentation and sex determination. Transducin-like enhancer pr... |
| Clonality                  | Monoclonal  |
| Concentration              | 0.1 mg/mL   |
| Clone Designation          | [TLE1/2062]   |

|                   |  |
|-------------------|--|
| Molecular Weight  | 83 kDa   |
| UniProt           | <a href="#">Q04724</a>   |
| Buffer            | PBS, 0.1% rBSA, 0.05% azide  |
| Source            | Animal   |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |