

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **MyoD1 (Rhabdomyosarcoma Marker) (MYOD1/2075R), CF740 conjugate, 0.1mg/mL, Clone: [MYOD1/2075R], CF 740, Rabbit, Monoclonal BOT-BNC742075-500**

|                            |  |
|----------------------------|--|
| Article Name               | MyoD1 (Rhabdomyosarcoma Marker) (MYOD1/2075R), CF740 conjugate, 0.1mg/mL, Clone: [MYOD1/2075R], CF 740, Rabbit, Monoclonal   |
| Biozol Catalog Number      | BOT-BNC742075-500  |
| Supplier Catalog Number    | BNC742075-500  |
| Alternative Catalog Number | BOT-BNC742075-500-500UL  |
| Manufacturer               | Biotium  |
| Host                       | Rabbit   |
| Category                   | Antikörper   |
| Application                | IHC  |
| Species Reactivity         | Human  |
| Immunogen                  | Recombinant full-length human MyoD1 protein  |
| Conjugation                | CF 740   |
| Product Description        | This antibody recognizes a phosphoprotein of 45 kDa, identified as MyoD1. This MAbs does not cross react with myogenin, Myf5, or Myf6. Antibody to MyoD1 labels the nuclei of myoblasts in developing muscle tissues. MyoD1 is not detected in normal adult... |
| Clonality                  | Monoclonal   |
| Concentration              | 0.1 mg/mL  |
| Clone Designation          | [MYOD1/2075R]  |

|                   |  |
|-------------------|--|
| Molecular Weight  | 45 kDa   |
| UniProt           | <a href="#">P15172</a>   |
| Buffer            | PBS, 0.1% rBSA, 0.05% azide  |
| Source            | Animal   |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |